


A Eukaryote without a Mitochondrial Organelle

Graphical Abstract



Authors

Anna Karnkowska, Vojtěch Vacek, Zuzana Zubáčová, ..., Joel B. Dacks, Čestmír Vlček, Vladimír Hampel

Correspondence

ankarn@biol.uw.edu.pl (A.K.), vlada@natur.cuni.cz (V.H.)

In Brief

Karnkowska et al. overturn the paradigm that eukaryotes must have mitochondria. Their genomic investigation of the anaerobic microbial eukaryote *Monocercomonoides sp.* reveals a complete lack of mitochondrial organelle and functions including Fe-S cluster synthesis, which is carried out in the cytosol by a laterally acquired bacterial pathway.

Highlights

- *Monocercomonoides sp.* is a eukaryotic microorganism with no mitochondria
- The complete absence of mitochondria is a secondary loss, not an ancestral feature
- The essential mitochondrial ISC pathway was replaced by a bacterial SUF system



A Eukaryote without a Mitochondrial Organelle

Anna Karnkowska,^{1,2,7,*} Vojtěch Vacek,¹ Zuzana Zubáčová,¹ Sebastian C. Treitli,¹ Romana Petrželková,³ Laura Eme,⁴ Lukáš Novák,¹ Vojtěch Žárský,¹ Lael D. Barlow,⁵ Emily K. Herman,⁵ Petr Soukal,¹ Miluše Hroudová,⁶ Pavel Doležal,¹ Courtney W. Stairs,⁴ Andrew J. Roger,⁴ Marek Eliáš,³ Joel B. Dacks,⁵ Čestmír Vlček,⁶ and Vladimír Hampl^{1,*}

¹Department of Parasitology, Charles University in Prague, Prague 12843, Czech Republic

²Department of Molecular Phylogenetics and Evolution, University of Warsaw, Warsaw 00478, Poland

³Department of Biology and Ecology, University of Ostrava, Ostrava 710 00, Czech Republic

⁴Department of Biochemistry and Molecular Biology, Dalhousie University, Halifax, NS B3H 4R2, Canada

⁵Department of Cell Biology, University of Alberta, Edmonton, AB T6G 2H7, Canada

⁶Institute of Molecular Genetics, Academy of Sciences of the Czech Republic, Prague 14220, Czech Republic

⁷Present address: Department of Botany, University of British Columbia, Vancouver, BC V6T 1Z4, Canada

*Correspondence: ankarn@biol.uw.edu.pl (A.K.), vlada@natur.cuni.cz (V.H.)

<http://dx.doi.org/10.1016/j.cub.2016.03.053>

SUMMARY

The presence of mitochondria and related organelles in every studied eukaryote supports the view that mitochondria are essential cellular components. Here, we report the genome sequence of a microbial eukaryote, the oxymonad *Monocercomonoides* sp., which revealed that this organism lacks all hallmark mitochondrial proteins. Crucially, the mitochondrial iron-sulfur cluster assembly pathway, thought to be conserved in virtually all eukaryotic cells, has been replaced by a cytosolic sulfur mobilization system (SUF) acquired by lateral gene transfer from bacteria. In the context of eukaryotic phylogeny, our data suggest that *Monocercomonoides* is not primitively amitochondrial but has lost the mitochondrion secondarily. This is the first example of a eukaryote lacking any form of a mitochondrion, demonstrating that this organelle is not absolutely essential for the viability of a eukaryotic cell.

INTRODUCTION

Mitochondria are organelles that arose through the endosymbiotic integration of an α -proteobacterial endosymbiont into the proto-eukaryote host cell. During the course of eukaryotic evolution, the genome and proteome of the mitochondrial compartment have been significantly modified, and many functions have been gained, lost, or relocated [1]. In extreme cases, the derivatives of mitochondria in anaerobic protists had become so modified that they had been overlooked [2] or not recognized as homologous to the mitochondrion [3]. Indeed, in the 1980s, the Archezoa hypothesis [4] proposed that some microbial eukaryotes primitively lacked mitochondria, peroxisomes, stacked Golgi apparatus, spliceosomal introns, and sexual reproduction. However, over the following decade, double-membraned organelles were identified in all investigated putative Archezoa. The final nail in the coffin of the Archezoa hypothesis was the demonstration that these organelles all contain some mitochondrial marker proteins, such as those involved in the iron-sulfur cluster

(ISC) Fe-S clusters biogenesis system, translocases, maturases, and/or molecular chaperones known to facilitate the import of proteins into mitochondria. It is now widely accepted that mitochondria or mitochondrion-related organelles (MROs) are essential compartments in all contemporary eukaryotes and that mitochondrial endosymbiosis took place before radiation of all extant eukaryotes [5].

Metamonada, originally part of the Archezoa, are now classified as one of the main clades of the eukaryotic “super-group” Excavata [6] and are comprised of microaerophilic or anaerobic unicellular eukaryotes that are often specialized parasites or symbionts. Detailed cell and molecular biological studies, including genome sequencing, have been undertaken only for three parasitic species from two metamonad lineages—*Giardia intestinalis* [7] and *Spiroplasma salmonicida* [8] (Fornicata) and *Trichomonas vaginalis* [9] (Parabasalia), which have provided important information regarding the functions of their MROs. The third lineage of metamonads, Preaxostyla, contains the basal paraphyletic free-living trimastixids and the derived endobiotic oxymonads [10]. The presence of mitochondrial homologs has been convincingly demonstrated in *Paratrimastix* (formerly *Trimastix*) *pyriformis*, although the biochemical functions of these organelles are largely unknown [11]. Endobiotic oxymonads belong to the least-studied former Archezoa. Here, we describe the first complete genome sequence analysis of an oxymonad, *Monocercomonoides* sp. PA203. We find that although this organism is a standard eukaryotic cell in other respects, it completely lacks any traces of a mitochondrion.

RESULTS AND DISCUSSION

Genome Characteristics

Using the 454 whole-genome shotgun sequencing methodology, we generated a draft genome sequence of the oxymonad *Monocercomonoides* sp. PA203, assembled into 2,095 scaffolds at $\sim 35\times$ coverage (see [Experimental Procedures](#)). The estimated size of the genome (~ 75 Mb) and the number of predicted protein-coding genes (16,629) is intermediate between what is found in diplomonads and *T. vaginalis* (Table 1). Almost 67% of predicted protein-coding genes contain introns (~ 1.9 introns per gene on average; Table 1). The assembly contains genes encoding tRNAs for all 20 amino acids, and ~ 50 ribosomal

Table 1. Overview of Metamonada Genomes

Taxa	Size (Mbp)	Guanine-Cytosine Content (%)	Protein-Coding Loci	Repetitive Regions	No. of Introns
<i>Monocercomonoides</i> sp. PA 203	~75	36.8	16,629	~38%	32,328
<i>Trichomonas vaginalis</i> isolate G3 [9]	~160	32.7	~60,000	~65%	65
<i>Giardia intestinalis</i> WB-C6 [7]	~11.7	49	6,480	9%	4
<i>Spironucleus salmonicida</i> ATCC 50377 [8]	12.9	33.4	8,076	5.2%	3

See also [Tables S1](#) and [S3](#).

DNA units were identified on small contigs outside the main assembly (see [Supplemental Experimental Procedures](#)). To estimate completeness of the genome sequence, we performed transcriptome mapping, in which 96.9% of transcripts mapped to the genome (see [Supplemental Experimental Procedures](#)), and checked the representation of core eukaryotic genes. Using the Core Eukaryotic Genes Mapping Approach (CEGMA) [12], we recovered 63.3% of core eukaryotic genes, a greater fraction than in the *G. intestinalis* genome (46.6%). However, when we excluded genes encoding mitochondrial proteins from the CEGMA dataset and used manually curated *Monocercomonoides* sp. gene models, the percentage of recovered genes increased to 90% ([Table S1](#)). For another set of 163 conserved eukaryotic genes used for phylogenomic analyses, the percentage of recovered genes exceeded 95% ([Table S2](#)). As the last measure of completeness, we identified 77 out of 78 conserved families of cytosolic eukaryotic ribosomal proteins [13] ([Table S3](#)), with the single exception of L41e, which is very short, difficult to detect, and has not been identified in other Metamonada genomes. Phylogenomic analysis ([Figure 1](#)) confirmed the relationship of *Monocercomonoides* sp. to *P. pyriformis* and other Metamonada and demonstrated that the *Monocercomonoides* lineage forms a much shorter branch relative to parabasalids and diplomonads. All these measures suggest that the assembled *Monocercomonoides* sp. genome sequence is nearly complete and its encoded proteins are, on average, less divergent than those of *G. intestinalis* and *T. vaginalis*.

With the first oxymonad genome sequence in hand, we focused our attention on one of the most puzzling aspects of their biology—the elusive nature of their mitochondrion.

Absence of Mitochondrial Proteins

No genes that are typically encoded on mitochondrial genomes (mtDNA) of other eukaryotes were found among the assembled scaffolds, suggesting that, like other metamonads, *Monocercomonoides* sp. lacks mtDNA. Next, we searched for homologs of nuclear genome-encoded proteins typically associated with mitochondria or MROs in other eukaryotes. The homologous core of the protein import machinery is regarded as strong evidence for the common origin of all mitochondria [14, 15]. As such, the presence of components of the translocases of the outer membrane (TOM) and inner membrane (TIM), sorting and assembly machinery (SAM) complex, and mitochondrial molecular chaperones (Hsp70 and Cpn60) in hydrogenosomes, mitosomes, and other MROs demonstrates that these organelles are related to mitochondria [16, 17]. While we were able to identify homologs of cytosolic chaperonins in the *Monocercomonoides* sp. genomic sequence, we were unable to identify homo-

logs of any component of the mitochondrial import machinery ([Figure 2A](#); [Experimental Procedures](#); [Tables S3](#) and [S4](#)).

All MROs, with the exception of the *G. intestinalis* mitosome [18], are known to export or import ATP and other metabolites typically using transporters from the mitochondrial carrier family (MCF) or, in mitosomes of the microsporidian *Encephalitozoon cuniculi* [19], by the bacterial-type (NTT-like) nucleotide transporters. We did not identify in the *Monocercomonoides* sp. genome any homologs of genes encoding known mitochondrial metabolite transport proteins ([Figure 2A](#); [Table S4](#)).

Fe-S clusters are essential biological cofactors associated with many different proteins and are therefore synthesized de novo in every organism across the tree of life [20]. In eukaryotes, this is done mostly by the mitochondrial ISC assembly system and the cytosolic iron-sulfur assembly (CIA) system [21]. Analyses of the *Monocercomonoides* sp. genome revealed the presence of a CIA system but a complete lack of components of the ISC system ([Figure 2A](#); [Table S3](#); [Experimental Procedures](#)).

We could not identify either of two possible enzymes involved in the synthesis of cardiolipin, a phospholipid specific for energy-transducing membranes [22]. The majority of eukaryotes synthesize cardiolipins, and the process is localized to mitochondria, but a complete lack of cardiolipin has been experimentally shown for *G. intestinalis*, *T. vaginalis*, and *E. cuniculi* [22]. Furthermore, we could not identify any component of the endoplasmic reticulum (ER)-mitochondria encounter structure (ERMES; [Figure 2A](#)) [23].

We identified only two orthologs of the set of proteins predicted to localize to the mitochondrion-related compartment of the closely related *P. pyriformis* [11]: aspartate/ornithine carbamoyltransferase family protein and pyridine nucleotide transhydrogenase. Neither protein has an exclusively mitochondrial localization in eukaryotes [24, 25], and the *Monocercomonoides* sp. orthologs do not contain predicted mitochondrial targeting sequences.

To complement the targeted homology-based searches, we also performed an extensive search for putative homologs of known mitochondrial proteins using a pipeline based on the Mitominer database [26], which was enriched with identified mitochondrial proteins of diverse anaerobic eukaryotes with MROs ([Experimental Procedures](#)). The search recovered 76 *Monocercomonoides* sp. proteins as candidates for functions in a putative mitochondrion ([Figure 2B](#); [Table S5](#)). Similarly to *G. intestinalis*, *T. vaginalis*, and *E. histolytica*, used as controls, the selected candidates were mainly proteins that are obviously not mitochondrial (e.g., histones) or for which the annotation is too general (e.g. “kinase domain-containing protein”), indicating that the specificity of the pipeline in organisms with




Figure 1. Unrooted Phylogeny of Eukaryotes Inferred from a 163-Protein Supermatrix

The tree displayed was inferred using PhyloBayes (CAT + Poisson substitution model). A maximum-likelihood (ML) tree inferred from the same supermatrix using RAxML (not shown) was very similar to the PhyloBayes tree, with the topological differences in the poorly resolved area comprising Chloroplastida, Cryptophyta, Glaucophyta, and Haptophyta, and in the position of Metamonada, in the ML tree placed sister (with strong bootstrap support) to Discoba. The branch support values shown are posterior probabilities (>0.95) from the PhyloBayes analysis and bootstrap values (>50%) from the ML analysis. Three branches are shown shortened to the indicated percentage of their actual length to fit them on the page. See also Table S2.

divergent mitochondrion is low. However, unlike all other control organisms, in which the search always recovered at least a few mitochondrial hallmark proteins, the set of 76 *Monocercomonoides* sp. candidates did not contain any such proteins. Only 11 of the *Monocercomonoides* candidates fall in the GO category “metabolism,” but they do not assemble any obvious metabolic pathway. In summary, this approach (Table S5) failed to reveal any credible set of mitochondrial protein in *Monocercomonoides* sp.

As an alternative to homology searches, we have also attempted to identify mitochondrial proteins by searching for several types of signature sequences. The matrix proteins of mitochondria and MROs are expected to contain conserved N-terminal targeting signals needed for the targeted import into MROs

[14]. We performed in silico prediction of mitochondrial targeting signals in the predicted *Monocercomonoides* sp. proteome and identified 107 candidate proteins (Figure 2A; Experimental Procedures; Table S6A). The presence of a predicted targeting signal by itself does not prove the targeting, as such amino acid sequences can also appear at random [27]. Functional annotation revealed that a majority of proteins recovered by this search fall into the Kyoto Encyclopedia of Genes and Genomes (KEGG) category “genetic information processing.” Given the absence of a mitochondrial genome, or organellar translation machinery, it is unlikely that these proteins function in an MRO. Only eight candidates were assigned to the KEGG category “metabolism,” and they are part of several different metabolic pathways. Finally, only three proteins were predicted to have a mitochondrial




Figure 2. Search Strategies for Proteins Functionally Related to the Mitochondrion in *Monocercomonoides*

(A) Search strategies for mitochondrial proteins and for protein-localization signatures in a canonical eukaryotic cell (details are given in [Supplemental Experimental Procedures](#)): (1) mitochondrial outer membrane (MOM)-targeted tail-anchored (TA) proteins ([Table S6B](#)), (2) proteins with a mitochondrial targeting signal ([Table S6A](#)), (3) β -barrel MOM proteins, (4) 41 mitochondrial hallmark proteins ([Table S4](#)), components of TOM and TIM translocases, cpn60, ERMES complex, ISC pathway components, cardiolipin synthase (CL).

(B) Semiautomatic pipeline for retrieving homologs of mitochondrial proteins from proteomes. We used a custom database for homology searching of mitochondrial proteins in the predicted proteomes of *Monocercomonoides* sp., *Giardia intestinalis*, *Entamoeba histolytica*, *Trichomonas vaginalis*, *Blastocystis* sp. subtype 7, and *Saccharomyces cerevisiae* ([Table S5](#)).

See also [Tables S4](#), [S5](#), and [S6](#).

targeting signal and homology to a Mitominer protein (hydrolyase-like family protein MONOS_10795, cytosolic TCP-1/cpn60 chaperonin family protein MONOS_13132, and ribonuclease

Z MONOS_6181). This also suggests that both pipelines failed to recover specific sets of mitochondrial proteins but instead detected only low-specificity “noise.”

The outer mitochondrial membranes accommodate two special classes of proteins, β -barrel and tail-anchored (TA) proteins, which are devoid of the N-terminal targeting signals and instead use specific C-terminal signals [28, 29]. We have identified 32 candidates for TA proteins in the predicted proteome, several of which appeared to be ER-targeted proteins. None of these had the hallmark characteristics of proteins targeted to the mitochondrial outer membrane (Figure 2A; Experimental Procedures; Table S6B). We also failed to identify any credible candidates for β -barrel outer membrane proteins (BOMPs) (Figure 2A; Experimental Procedures).

In summary, our comprehensive examination of the *Monocercomonoides* sp. genome based on homology searches and searches for specific N-terminal and C-terminal signals failed to recover proteins typically associated with MROs, including mitochondrial translocases, metabolite transporters and the ISC system for Fe-S cluster synthesis, ERMES, and enzymes responsible for cardiolipin synthesis.

In order to verify that our inability to find any reliable mitochondrial proteins is not caused by possible unprecedented divergence of *Monocercomonoides* sp. proteins or a failure of our methods, we searched for hallmark proteins of another cellular system, so far not observed in *Monocercomonoides* sp.—the Golgi complex. In this case, using homology-based searches, we detected numerous Golgi-associated proteins, including components of the COPI, AP-1, AP-3, AP-4, COG, GARP, TRAPPI, and Retromer complexes and Rab GTPases regulating transport to and from the Golgi (Table S3). This suggests the presence of Golgi-like compartments in oxymonads [30], despite the absence of a cytologically discernible Golgi apparatus.

The specific absence of mitochondria-associated proteins in *Monocercomonoides* sp. implies the legitimate absence of a mitochondrial compartment. If so, then how does the *Monocercomonoides* cell function without this organelle?

Energy Metabolism without a Mitochondrion

In order to compare the metabolism of *Monocercomonoides* sp. with anaerobic protists retaining mitochondrial compartments, we performed manual annotation of proteins of core pathways of energy metabolism normally associated with the presence and function of a MRO. As with many other organisms with secondarily reduced mitochondria, the *Monocercomonoides* sp. genome does not encode any enzymes for aerobic energy generation (e.g., TCA cycle or electron transport chain proteins). We did identify a complete set of glycolytic enzymes, including the alternative enzymes for anaerobic glycolysis [31], as well as the anaerobic fermentation enzymes pyruvate:ferredoxin oxidoreductase (PFOR) and [FeFe]-hydrogenases (Table S3). [FeFe]-hydrogenase maturases were absent, which is not unprecedented as they are also missing from *G. intestinalis* and *E. histolytica*, anaerobic parasites that are both capable of cytosolic H_2 production [32, 33]. Neither PFOR nor [FeFe]-hydrogenase has a predicted mitochondrial targeting sequence, and heterologous expression in *T. vaginalis* suggests a cytosolic localization of PFOR (Figure S1). In summary, *Monocercomonoides* sp. glucose metabolism appears to produce ATP via substrate-level phosphorylation steps in an extended glycolysis pathway, and the reduced co-factors are re-oxidized by fermentation, ultimately producing acetate and ethanol, or by [FeFe]-hy-

drogenase producing hydrogen gas. The situation in *Monocercomonoides* sp. is virtually identical to *G. intestinalis* and *E. histolytica*, which independently reduced their mitochondria to mitosomes and all the ATP production occurs in the cytosol [34–36].


In addition to extended glycolysis, *Monocercomonoides* sp. contains a complete set of three genes for enzymes involved in arginine deiminase pathway—arginine deiminase, ornithine carbamoyltransferase, and carbamate kinase. This pathway may also be used for ATP production by arginine degradation as in *T. vaginalis* and *G. intestinalis* [37, 38]. In *G. intestinalis*, this pathway produces eight times more ATP than sugar metabolism.

Fe-S Cluster Assembly without a Mitochondrion

Every eukaryotic cell contains a CIA machinery, which assists the final stages of the assembly of Fe-S clusters in proteins functioning in the eukaryotic cytosol and nucleus. Eight proteins were shown to be involved in the CIA pathway in yeast and humans: Cfd1, NUBP1 (Nbp35), NARFL (Nar1), CIAO1 (Cia1), Dre2, Tah18, Cia2, and MMS19. Four of them (i.e., Nbp35, Nar1, Cia1, and Cia2) [21] are conserved among eukaryotes and also present in the *Monocercomonoides* sp. genome (Table S3). We did not identify Cfd1 and MMS19, which are missing from many other eukaryotes, and Dre2 and Tah18, which are missing from the anaerobic protists containing MROs (including *E. histolytica*, *Mastigamoeba balamuthi*, *T. vaginalis*, *G. intestinalis*, and *Blastocystis* sp.) [21].

Despite the presence of the CIA pathway, it is commonly suggested that mitochondria and related organelles are essential to eukaryotic cells because the mitochondrial ISC system plays a critical role in the initial phase of the formation of cytosolic Fe-S clusters [20]. Although the ISC system is a near-universally conserved pathway in eukaryotes and seems to be the unifying feature of mitochondria and related organelles, genes encoding proteins of the mitochondrial ISC pathway have not been detected in the *Monocercomonoides* sp. genome. The functional replacement of the ISC system has been reported for only two lineages, *Pygsuia biforma* (Breviatea) and Archamoebae. A methanoarchaeal sulfur mobilization (SUF) system [39] or a bacterial nitrogen fixation (NIF) [40] has apparently replaced the ISC system in the *P. biforma* and the Archamoebae lineages, respectively. Conflicting data exist on the localization of the NIF system in *E. histolytica* [41, 42]; however, in *M. balamuthi*, the NIF system localizes in the cytosol and the MRO [43].

The major issue remains: how does *Monocercomonoides* sp. form Fe-S clusters? Unexpectedly, we identified genes encoding four subunits of the SUF system: SufB, SufC, and fused SufS and SufU (Table S3). SufS is a “two-component” cysteine desulfurase, and its activity might be enhanced by SufE or SufU [44, 45]. In *Monocercomonoides* sp., SufS is fused with SufU, which is a unique feature. SufB and SufC can form a scaffold complex in prokaryotes, and SufB2C2 complex is capable of binding and transferring 4Fe-4S clusters to a recipient apoprotein [46]. All identified SUF system proteins apparently retain all important catalytic sites (Figure S2) and may perform de novo Fe-S clusters biogenesis by themselves or in concert with the CIA machinery. The SUF system for Fe-S cluster synthesis is found in plastids, bacteria, and archaea and has also been found in two microbial eukaryotes *P. biforma* [39] and *Blastocystis* sp. [47]. The



(legend on next page)




Figure 4. Reductive Evolution of Mitochondria in Metamonads

Transition to an anaerobic lifestyle occurred in a common ancestor of metamonads and was followed by reduction of mitochondria to MROs, accompanied by the loss of cristae and genome, and the transition to anaerobic metabolism. The ISC pathway for Fe-S cluster synthesis was present in a metamonad common ancestor. Further reduction to a mitosome took place in the *Giardia intestinalis* lineage. We propose that in the common ancestor of *Paratrimastix pyriformis* and *Monocercomonoides*, a Suf system acquired through LGT from bacteria substituted the MRO-localized ISC system. Subsequently, the MRO was lost completely in the lineage leading to *Monocercomonoides* sp. Localization of the Suf pathway in *P. pyriformis* is unknown.

presence of spliceosomal introns in the putative SufC and SufSU of *Monocercomonoides* confirms that these proteins are not prokaryotic contamination. Furthermore, fluorescence in situ hybridization (FISH) with *sufB* and *sufC* gene probes demonstrated their presence in the *Monocercomonoides* sp. nucleus (Figure S3). Importantly, homologs of these proteins were detected in the *P. pyriformis* genome, the closest sequenced relative to *Monocercomonoides*. The Suf system components of both *Monocercomonoides* sp. and *P. pyriformis* do not contain recognizable mitochondrial targeting signals, and our experiments with heterologous expression of *Monocercomonoides* sp. SufB and SufC proteins in *T. vaginalis* (Figure 3A) and SufC protein in yeast (Figure 3B) support a cytosolic localization. Phylogenetic analyses indicate that this Suf system was acquired by an ancestor of *Monocercomonoides* and *Paratrimastix* by lateral gene transfer (LGT) from bacteria independently of all other Suf-containing eukaryotes (Figure 3C). We propose that the acquisition of a cytosolic Suf system made the ancestral ISC system in the mitochondrion dispensable, which led to its loss

and, in the *Monocercomonoides* lineage, to the complete loss of MROs (Figure 4).

Conclusions

Mitochondria and related organelles are currently considered to be indispensable components of eukaryotic cells. The genome sequence of *Monocercomonoides* sp. reported here suggests that this is not the case. Despite extensive searches, no mitochondrial marker proteins such as membrane protein translocases and metabolite transporters were identified. Crucially, the mitochondrion-specific ISC pathway for Fe-S cluster biogenesis is absent and apparently was replaced by a bacterial Suf system that functions in the cytosol. On the other hand, genes encoding other features once thought to be absent from these divergent eukaryotic cells, i.e., the Golgi body, were readily identifiable. The genome also contains genes for essential cytosolic pathways of energy metabolism, although we did observe examples of metabolic streamlining characteristic of other anaerobic or microaerophilic eukaryotes.

Figure 3. Heterologous Expression of *Monocercomonoides* sp. Suf System Proteins and Phylogeny of Concatenated SufB, SufC, and SufS Homologs

(A) Heterologous expression of *Monocercomonoides* sp. SufB and SufC proteins in *Trichomonas vaginalis*. *Monocercomonoides* sp. proteins with a C-terminal HA tag were expressed in *T. vaginalis* and visualized by an anti-HA antibody (green). The signal of the anti-HA antibody does not co-localize with hydrogenosomes stained using an anti-malic enzyme antibody (red). The nucleus was stained using DAPI (blue). Scale bar, 10 μ m.

(B) Heterologous expression of *Monocercomonoides* sp. SufC protein in *Saccharomyces cerevisiae*. *Monocercomonoides* sp. proteins tagged with GFP were expressed in *S. cerevisiae* (green). The GFP signal does not co-localize with the yeast mitochondria stained by Mitotracker (red). Scale bar, 10 μ m.

(C) Unrooted ML tree of concatenated SufB, SufC, and SufS sequences. Bootstrap support values above 50 and posterior probabilities greater than 0.75 are shown. *Monocercomonoides* sp. and *Paratrimastix pyriformis* are shown in red, eukaryotic plastids and cyanobacteria in green, *Blastocystis* sp. and *Pygusua biforma* in orange, bacteria in gray, and archaea in blue.

See also Figures S1–S3.

Reduction of mitochondria is known from various eukaryotic lineages adapted to anaerobic lifestyle [48]. Mitosomes in *Giardia*, *Entamoeba*, and Microsporidia represent the most extreme cases of mitochondrial reduction known to date, and yet they still contain recognizable mitochondrial protein translocases and usually an ISC system. The specific absence of all these mitochondrial proteins in the genome of *Monocercomonoides* sp. indicates that this eukaryote has dispensed with the mitochondrial compartment completely. In principle, we cannot exclude the possibility that a mitochondrion exists in *Monocercomonoides* sp. whose protein composition has been altered entirely. However, such a hypothetical organelle could not be recognized as a mitochondrion homolog by any available means. Without any positive evidence for the latter scenario, we suggest that the complete absence of mitochondrial markers and pathways points to the bona fide absence of the organelle. Because all known oxymonads are obligate animal symbionts, and mitochondrial homologs are present in the close free-living sister lineage *Paratrimastix*, the absence of mitochondrion in *Monocercomonoides* sp. must be secondary. We hypothesize that the acquisition of the SUF system predated the loss of the mitochondrial ISC system in the common ancestor of Preaxostyla and allowed for the complete loss of the organelle in *Monocercomonoides* sp. lineage, the first known truly secondarily amitochondriate eukaryote.

EXPERIMENTAL PROCEDURES

Genome and Transcriptome Sequencing

All experiments were performed on the *Monocercomonoides* sp. PA203 strain. The culture (2 L with a cell density of approximately 4×10^5 cells/mL) was filtered to remove most of the bacteria before isolation of DNA (culturing and filtration details in [Supplemental Experimental Procedures](#)). DNA was isolated using DNeasy Blood and Tissue Kit (QIAGEN). Total genomic DNA was sequenced using a Genome Sequencer 454 GS FLX+ with XL+ reagents. A total of seven sequencing runs were performed, including four shotgun runs on libraries with the average fragment length of 500 to 800 and three runs on a 3-kb paired-end library. Two RNA sequencing (RNA-seq) experiments were performed using 454 and Illumina sequencing platforms. Details of sequencing are given in [Supplemental Experimental Procedures](#).

Roche's assembler Newbler v.2.6 was used to generate a genome sequence assembly from 454 single and pair end reads. The final assembly consisted of 2,095 scaffolds spanning almost 75 Mb of the genome. The N50 scaffold size is 71.4 kb. Transcriptome assembly of the 454 data was performed by Newbler v.2.8 with default parameters, and Illumina-generated transcriptomic data were assembled using Trinity [49] (details in [Supplemental Experimental Procedures](#)). The CEGMA [12] was used to estimate the number of conserved eukaryotic genes in the *Monocercomonoides* sp. genome assembly (Table S1) and presence of cytosolic ribosomal eukaryotic proteins as an additional measure of completeness (Table S3).

Genome Annotation and Gene Searching

For the structural annotation, Augustus v.2.7 [50, 51], PASA2 [52], and EVM [53] were used. Gene models of particular interest were manually evaluated with the help of RNA-seq data or considering conservation with homologs (details in [Supplemental Experimental Procedures](#)).

Functional annotation was assigned to genes by similarity searches of predicted proteins using BLASTP [54] against the NCBI non-redundant protein database [55] and HMMER3 [56] searches of domain hits in the Pfam protein families database [57]. Additional annotation was performed using the KEGG automatic annotation server [58]. Annotation files are available at the web page <http://www.protistologie.cz/hamp/lab/data.html>.

tRNA genes were predicted with tRNAscan-SE [59]; rDNA sequences were not present in the original main assembly, but they were identified in contigs not assembled into scaffolds and added to the main assembly.

The *Monocercomonoides* sp. genome database was searched using the TBLASTN [54] algorithm, and *Monocercomonoides* proteome database and six-frame translation of the genomic sequence were searched using the BLASTP [54] algorithm or the profile hidden Markov model (HMM) searching method *phmmer* from the HMMER3 [56] package. We used a wide range of queries described in [Supplemental Experimental Procedures](#).

Phylogenetic Analyses

We performed a number of maximum-likelihood and Bayesian phylogenetic analyses: (1) phylogenomic analyses of eukaryotes based on 163 genes and 70 taxa; (2) phylogenetic analyses of genes for SUF pathway enzymes; and (3) individual gene trees to support functional annotation of genes (details in [Supplemental Experimental Procedures](#)).

Subcellular Localization Prediction

Subcellular localization prediction for the *Monocercomonoides* sp. proteome was performed using TargetP v.1.1 [60] and MitoProt II v.1.101 [61]. TA proteins were identified and analyzed based on presence of a transmembrane domain (TMD) of moderate hydrophobicity flanked by positively charged residues [29, 62] (details in [Supplemental Experimental Procedures](#)). BOMPs were identified based on the presence of a conserved C-terminal β -signal, using a previously described pipeline [63].

Mitochondrial Protein Searching Using a Mitominer-Based Database

We prepared a custom database of mitochondrial proteins to search for genes encoding proteins with putative mitochondrial localization. The custom database was based on the MitoMiner database [26] reference set containing 12,925 proteins from 11 eukaryotic mitochondrial proteomes, which was enriched by known or predicted MRO-localized proteins of *E. histolytica*, *G. intestinalis*, *P. biforma*, *S. salmonicida*, *T. vaginalis*, and *P. pyriformis*. Homologs of proteins from this database were searched in the predicted proteome of *Monocercomonoides* sp. and in the predicted proteomes of *Blastocystis* sp., *E. histolytica*, *G. intestinalis*, *S. cerevisiae*, and *T. vaginalis*, which were used as control datasets. While searching the control datasets, the proteins of the searched organism were removed from the custom database. In the last step, only those candidates were kept whose first hit in the NCBI database [55] contained a predictable mitochondrial targeting signal (score > 0.5 in TargetP v.1.1 [60] and MitoProt II v.1.101 [61]). Further details are given in [Supplemental Experimental Procedures](#).

FISH

We performed FISH experiments with labeled probes to determine whether the genes for SUF system proteins physically reside in the *Monocercomonoides* sp. genome or represent bacterial contamination. Details on preparation of labeled probes are given in [Supplemental Experimental Procedures](#).

One liter of *Monocercomonoides* sp. culture was filtered to remove bacteria, and the cells were pelleted by centrifugation for 10 min at $2,000 \times g$ at 4°C . FISH with digoxigenin-labeled probes was performed according to a previously described procedure [64] omitting the colchicine procedure. Cell nuclei and the probes were denatured under a coverslip in a single step in $50 \mu\text{L}$ of 50% formamide in $2 \times \text{SSC}$ at 70°C for 5 min. Preparations were observed using an IX81 microscope (Olympus) equipped with an IX2-UCB camera. Images were processed using Cell software (Olympus) and ImageJ 1.42q.

Heterologous Protein Expression and Microscopy in *Trichomonas vaginalis*

The *T. vaginalis* transfection system was used to assess subcellular localization of SufB, SufC, and PFOR proteins. *Monocercomonoides* sp. cDNA preparation was performed as described for transcriptome sequencing ([Supplemental Experimental Procedure](#)). Constructs with the hemagglutinin (HA) tag fused to the 3' end of the coding sequences of the studied genes were prepared and expressed in *T. vaginalis*, an anaerobic protist related to *Monocercomonoides* sp. and bearing a hydrogenosome (details are given in [Supplemental Experimental Procedures](#)). *Monocercomonoides* sp. proteins

expressed in *T. vaginalis* cells were visualized using standard techniques [14] (details are given in [Supplemental Experimental Procedures](#)).

Saccharomyces cerevisiae Heterologous Expression System

This expression system was used to confirm the results from the *T. vaginalis* expression system for SufC protein. The procedure was analogous to the one described in [11]. Details are given in [Supplemental Experimental Procedures](#).

ACCESSION NUMBERS

Sequence data for the genome reads (experiment number SRX1470187), the 454 transcriptome reads sequenced using the 454 platform (experiment number SRX1453820), and the Illumina transcriptome reads sequenced using the Illumina platform (experiment number SRX1453675) have been deposited to the NCBI Sequence Read Archive under accession number SRA: SRP066769. The accession number for the *Monocercomonoides* sp. PA203 genome reported in this paper is GenBank: LSR000000000. The accession number for the 454 transcriptome project reported in this paper is GenBank: GEEG000000000. The accession number for the Illumina transcriptome project reported in this paper is GenBank: GEEL000000000. The versions described in this paper are versions LSR010000000, GEEG010000000, and GEEL010000000. Further additional information on the genome analysis can be found at <http://www.protistologie.cz/hampllab/data.html>.

SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures, three figures, and six tables and can be found with this article online at <http://dx.doi.org/10.1016/j.cub.2016.03.053>.

AUTHOR CONTRIBUTIONS

The project was conceived in the laboratory of V.H. with the contribution of J.B.D. Genome and 454 transcriptome sequencing was performed by the Laboratory of Genomics and Bioinformatics. A.K. and V.H. coordinated the project. Z.Z. isolated genomic DNA. V.V. and V.H. isolated RNA. M.H. prepared sequencing libraries. Č.V. and A.K. assembled data. A.K. curated data, analyzed genomic and transcriptomic data, and conducted gene prediction and automatic functional annotation. A.K., V.V., S.C.T., R.P., L.N., V.Ž., L.D.B., E.K.H., M.E., and V.H. performed manual annotation. A.K., V.V., P.D., C.W.S., and V.H. performed mitochondrial gene searching. Z.Z. performed FISH experiments. V.V., Z.Z., and S.C.T. performed immunolocalization experiments. A.K., V.V., R.P., L.D.B., E.K.H., P.S., and L.E. performed phylogenetic analyses. A.K., V.V., Z.Z., S.C.T., and R.P. prepared figures. A.K. and V.H. wrote the manuscript in collaboration with A.J.R., M.E., and J.B.D., and all authors edited and approved the manuscript.

ACKNOWLEDGMENTS

A.K. and V.H. were supported by the Ministry of Education, Youth and Sports of CR within the National Sustainability Program II (Project BIOCEV-FAR) LQ1604 and by the project “BIOCEV” (CZ.1.05/1.1.00/02.0109). V.H. and sequencing were supported by Czech Science foundation project P506-12-1010. Z.Z. and localization experiments were funded by Czech Science foundation project 510 13-22333P. E.K.H. was supported by a Vanier Canada Graduate Scholarship and an Alberta Innovates – Health Solutions Graduate Studentship. The work of L.E., C.W.S., and A.J.R. was supported by a regional partnerships program grant (62809) from the Canadian Institute of Health Research and the Nova Scotia Health Research Foundation. The work of L.D.B., E.K.H., and J.B.D. was supported by an NSERC Discovery grant and an Alberta Innovates Technology Futures New Faculty Award to J.B.D. R.P. and M.E. were supported by Czech Science foundation project 15-16406S.

Received: December 23, 2015

Revised: March 5, 2016

Accepted: March 23, 2016

Published: May 12, 2016

REFERENCES

- Huynen, M.A., Duarte, I., and Szklarczyk, R. (2013). Loss, replacement and gain of proteins at the origin of the mitochondria. *Biochim. Biophys. Acta* 1827, 224–231.
- Tovar, J., León-Avila, G., Sánchez, L.B., Sutak, R., Tachezy, J., van der Giezen, M., Hernández, M., Müller, M., and Lucocq, J.M. (2003). Mitochondrial remnant organelles of Giardia function in iron-sulphur protein maturation. *Nature* 426, 172–176.
- Lindmark, D.G., and Müller, M. (1973). Hydrogenosome, a cytoplasmic organelle of the anaerobic flagellate *Tritrichomonas foetus*, and its role in pyruvate metabolism. *J. Biol. Chem.* 248, 7724–7728.
- Cavalier-Smith, T. (1987). Eukaryotes with no mitochondria. *Nature* 326, 332–333.
- Gray, M.W. (2012). Mitochondrial evolution. *Cold Spring Harb. Perspect. Biol.* 4, a011403.
- Adl, S.M., Simpson, A.G.B., Lane, C.E., Lukeš, J., Bass, D., Bowser, S.S., Brown, M.W., Burki, F., Dunthorn, M., Hampl, V., et al. (2012). The revised classification of eukaryotes. *J. Eukaryot. Microbiol.* 59, 429–493.
- Morrison, H.G., McArthur, A.G., Gillin, F.D., Aley, S.B., Adam, R.D., Olsen, G.J., Best, A.A., Cande, W.Z., Chen, F., Cipriano, M.J., et al. (2007). Genomic minimalism in the early diverging intestinal parasite *Giardia lamblia*. *Science* 317, 1921–1926.
- Xu, F., Jerlström-Hultqvist, J., Einarsson, E., Astvaldsson, A., Svärd, S.G., and Andersson, J.O. (2014). The genome of *Spironucleus salmonicida* highlights a fish pathogen adapted to fluctuating environments. *PLoS Genet.* 10, e1004053.
- Carlton, J.M., Hirt, R.P., Silva, J.C., Delcher, A.L., Schatz, M., Zhao, Q., Wortman, J.R., Bidwell, S.L., Alsmark, U.C.M., Besteiro, S., et al. (2007). Draft genome sequence of the sexually transmitted pathogen *Trichomonas vaginalis*. *Science* 315, 207–212.
- Zhang, Q., Táborský, P., Silberman, J.D., Pánek, T., Čepička, I., and Simpson, A.G.B. (2015). Marine isolates of *Trimastix marina* form a plesiomorphic deep-branching lineage within Preaxostyla, separate from other known Trimastigids (*Paratrimastix* n. gen.). *Protist* 166, 468–491.
- Zubáčová, Z., Novák, L., Bublíková, J., Vacek, V., Fousek, J., Rídl, J., Tachezy, J., Doležal, P., Vlček, C., and Hampl, V. (2013). The mitochondrion-like organelle of *Trimastix pyriformis* contains the complete glycine cleavage system. *PLoS ONE* 8, e55417.
- Parra, G., Bradnam, K., and Korf, I. (2007). CEGMA: a pipeline to accurately annotate core genes in eukaryotic genomes. *Bioinformatics* 23, 1061–1067.
- Lecompte, O., Ripp, R., Thierry, J.C., Moras, D., and Poch, O. (2002). Comparative analysis of ribosomal proteins in complete genomes: an example of reductive evolution at the domain scale. *Nucleic Acids Res.* 30, 5382–5390.
- Doležal, P., Likic, V., Tachezy, J., and Lithgow, T. (2006). Evolution of the molecular machines for protein import into mitochondria. *Science* 313, 314–318.
- Zarsky, V., Tachezy, J., and Doležal, P. (2012). Tom40 is likely common to all mitochondria. *Curr. Biol.* 22, R479–R481, author reply R481–R482.
- Doležal, P., Smid, O., Rada, P., Zubáčová, Z., Bursac, D., Suták, R., Nebesárová, J., Lithgow, T., and Tachezy, J. (2005). Giardia mitosomes and trichomonad hydrogenosomes share a common mode of protein targeting. *Proc. Natl. Acad. Sci. USA* 102, 10924–10929.
- Burri, L., Williams, B.A.P., Bursac, D., Lithgow, T., and Keeling, P.J. (2006). Microsporidian mitosomes retain elements of the general mitochondrial targeting system. *Proc. Natl. Acad. Sci. USA* 103, 15916–15920.
- Jedelský, P.L., Doležal, P., Rada, P., Pyrih, J., Smid, O., Hrdý, I., Sedínová, M., Marcínčiková, M., Voleman, L., Perry, A.J., et al. (2011). The minimal proteome in the reduced mitochondrion of the parasitic protist *Giardia intestinalis*. *PLoS ONE* 6, e17285.

19. Tsaousis, A.D., Kunji, E.R.S., Goldberg, A.V., Lucocq, J.M., Hirt, R.P., and Embley, T.M. (2008). A novel route for ATP acquisition by the remnant mitochondria of *Encephalitozoon cuniculi*. *Nature* *453*, 553–556.
20. Lill, R. (2009). Function and biogenesis of iron-sulphur proteins. *Nature* *460*, 831–838.
21. Tsaousis, A.D., Gentekaki, E., Eme, L., Gaston, D., and Roger, A.J. (2014). Evolution of the cytosolic iron-sulfur cluster assembly machinery in *Blastocystis* species and other microbial eukaryotes. *Eukaryot. Cell* *13*, 143–153.
22. Tian, H.-F., Feng, J.-M., and Wen, J.-F. (2012). The evolution of cardiolipin biosynthesis and maturation pathways and its implications for the evolution of eukaryotes. *BMC Evol. Biol.* *12*, 32.
23. Wideman, J.G., Gawryluk, R.M.R., Gray, M.W., and Dacks, J.B. (2013). The ancient and widespread nature of the ER-mitochondria encounter structure. *Mol. Biol. Evol.* *30*, 2044–2049.
24. Yarlett, N., Lindmark, D.G., Goldberg, B., Moharrami, M.A., and Bacchi, C.J. (1994). Subcellular localization of the enzymes of the arginine dihydrolase pathway in *Trichomonas vaginalis* and *Tritrichomonas foetus*. *J. Eukaryot. Microbiol.* *41*, 554–559.
25. Yousuf, M.A., Mi-ichi, F., Nakada-Tsukui, K., and Nozaki, T. (2010). Localization and targeting of an unusual pyridine nucleotide transhydrogenase in *Entamoeba histolytica*. *Eukaryot. Cell* *9*, 926–933.
26. Smith, A.C., Blackshaw, J.A., and Robinson, A.J. (2012). MitoMiner: a data warehouse for mitochondrial proteomics data. *Nucleic Acids Res.* *40*, D1160–D1167.
27. Lucattini, R., Likić, V.A., and Lithgow, T. (2004). Bacterial proteins predisposed for targeting to mitochondria. *Mol. Biol. Evol.* *21*, 652–658.
28. Denic, V. (2012). A portrait of the GET pathway as a surprisingly complicated young man. *Trends Biochem. Sci.* *37*, 411–417.
29. Borgese, N., Brambillasca, S., and Colombo, S. (2007). How tails guide tail-anchored proteins to their destinations. *Curr. Opin. Cell Biol.* *19*, 368–375.
30. Mowbray, K., and Dacks, J.B. (2009). Evolution and diversity of the Golgi body. *FEBS Lett.* *583*, 3738–3745.
31. Liapounova, N.A., Hampl, V., Gordon, P.M.K., Sensen, C.W., Gedamu, L., and Dacks, J.B. (2006). Reconstructing the mosaic glycolytic pathway of the anaerobic eukaryote *Monocercomonoides*. *Eukaryot. Cell* *5*, 2138–2146.
32. Lloyd, D., Ralphs, J.R., and Harris, J.C. (2002). *Giardia intestinalis*, a eukaryote without hydrogenosomes, produces hydrogen. *Microbiology* *148*, 727–733.
33. Nixon, J.E.J., Field, J., McArthur, A.G., Sogin, M.L., Yarlett, N., Loftus, B.J., and Samuelson, J. (2003). Iron-dependent hydrogenases of *Entamoeba histolytica* and *Giardia lamblia*: activity of the recombinant entamoebic enzyme and evidence for lateral gene transfer. *Biol. Bull.* *204*, 1–9.
34. van der Giezen, M., and Tovar, J. (2005). Degenerate mitochondria. *EMBO Rep.* *6*, 525–530.
35. Müller, M., Mentel, M., van Hellemond, J.J., Henze, K., Woehle, C., Gould, S.B., Yu, R.-Y., van der Giezen, M., Tielens, A.G.M., and Martin, W.F. (2012). Biochemistry and evolution of anaerobic energy metabolism in eukaryotes. *Microbiol. Mol. Biol. Rev.* *76*, 444–495.
36. Makiuchi, T., and Nozaki, T. (2014). Highly divergent mitochondrion-related organelles in anaerobic parasitic protozoa. *Biochimie* *100*, 3–17.
37. Schofield, P.J., Edwards, M.R., Matthews, J., and Wilson, J.R. (1992). The pathway of arginine catabolism in *Giardia intestinalis*. *Mol. Biochem. Parasitol.* *51*, 29–36.
38. Yarlett, N., Martinez, M.P., Moharrami, M.A., and Tachezy, J. (1996). The contribution of the arginine dihydrolase pathway to energy metabolism by *Trichomonas vaginalis*. *Mol. Biochem. Parasitol.* *78*, 117–125.
39. Stairs, C.W., Eme, L., Brown, M.W., Mutsaers, C., Susko, E., Delleire, G., Soanes, D.M., van der Giezen, M., and Roger, A.J. (2014). A SUF Fe-S cluster biogenesis system in the mitochondrion-related organelles of the anaerobic protist *Pygsuia*. *Curr. Biol.* *24*, 1176–1186.
40. van der Giezen, M., Cox, S., and Tovar, J. (2004). The iron-sulfur cluster assembly genes *iscS* and *iscU* of *Entamoeba histolytica* were acquired by horizontal gene transfer. *BMC Evol. Biol.* *4*, 7.
41. Maralíkova, B., Ali, V., Nakada-Tsukui, K., Nozaki, T., van der Giezen, M., Henze, K., and Tovar, J. (2010). Bacterial-type oxygen detoxification and iron-sulfur cluster assembly in amoebal relict mitochondria. *Cell. Microbiol.* *12*, 331–342.
42. Mi-ichi, F., Abu Yousuf, M., Nakada-Tsukui, K., and Nozaki, T. (2009). Mitosomes in *Entamoeba histolytica* contain a sulfate activation pathway. *Proc. Natl. Acad. Sci. USA* *106*, 21731–21736.
43. Nývltová, E., Šuták, R., Harant, K., Šedinová, M., Hrdý, I., Paces, J., Vlček, Č., and Tachezy, J. (2013). NIF-type iron-sulfur cluster assembly system is duplicated and distributed in the mitochondria and cytosol of *Mastigamoeba balamuthi*. *Proc. Natl. Acad. Sci. USA* *110*, 7371–7376.
44. Loiseau, L., Ollagnier-de-Choudens, S., Nachin, L., Fontecave, M., and Barras, F. (2003). Biogenesis of Fe-S cluster by the bacterial Suf system: SufS and SufE form a new type of cysteine desulfurase. *J. Biol. Chem.* *278*, 38352–38359.
45. Riboldi, G.P., de Oliveira, J.S., and Frazzon, J. (2011). Enterococcus faecalis SufU scaffold protein enhances SufS desulfurase activity by acquiring sulfur from its cysteine-153. *Biochim. Biophys. Acta* *1814*, 1910–1918.
46. Chahal, H.K., and Outten, F.W. (2012). Separate FeS scaffold and carrier functions for SufB₂C₂ and SufA during in vitro maturation of [2Fe2S] Fdx. *J. Inorg. Biochem.* *116*, 126–134.
47. Tsaousis, A.D., Ollagnier de Choudens, S., Gentekaki, E., Long, S., Gaston, D., Stechmann, A., Vinella, D., Py, B., Fontecave, M., Barras, F., et al. (2012). Evolution of Fe/S cluster biogenesis in the anaerobic parasite *Blastocystis*. *Proc. Natl. Acad. Sci. USA* *109*, 10426–10431.
48. Maguire, F., and Richards, T.A. (2014). Organelle evolution: a mosaic of ‘mitochondrial’ functions. *Curr. Biol.* *24*, R518–R520.
49. Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., et al. (2011). Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat. Biotechnol.* *29*, 644–652.
50. Stanke, M., and Waack, S. (2003). Gene prediction with a hidden Markov model and a new intron submodel. *Bioinformatics* *19* (Suppl 2), ii215–ii225.
51. Stanke, M., Schöffmann, O., Morgenstern, B., and Waack, S. (2006). Gene prediction in eukaryotes with a generalized hidden Markov model that uses hints from external sources. *BMC Bioinformatics* *7*, 62.
52. Haas, B.J., Delcher, A.L., Mount, S.M., Wortman, J.R., Smith, R.K., Jr., Hannick, L.I., Maiti, R., Ronning, C.M., Rusch, D.B., Town, C.D., et al. (2003). Improving the Arabidopsis genome annotation using maximal transcript alignment assemblies. *Nucleic Acids Res.* *31*, 5654–5666.
53. Haas, B.J., Salzberg, S.L., Zhu, W., Pertea, M., Allen, J.E., Orvis, J., White, O., Buell, C.R., and Wortman, J.R. (2008). Automated eukaryotic gene structure annotation using EvidenceModeler and the Program to Assemble Spliced Alignments. *Genome Biol.* *9*, R7.
54. Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D.J. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* *25*, 3389–3402.
55. Pruitt, K.D., Tatusova, T., and Maglott, D.R. (2005). NCBI Reference Sequence (RefSeq): a curated non-redundant sequence database of genomes, transcripts and proteins. *Nucleic Acids Res.* *33*, D501–D504.
56. Finn, R.D., Clements, J., and Eddy, S.R. (2011). HMMER web server: interactive sequence similarity searching. *Nucleic Acids Res.* *39*, W29–37.
57. Punta, M., Coghill, P.C., Eberhardt, R.Y., Mistry, J., Tate, J., Boursnell, C., Pang, N., Forslund, K., Ceric, G., Clements, J., et al. (2012). The Pfam protein families database. *Nucleic Acids Res.* *40*, D290–D301.
58. Moriyama, Y., Itoh, M., Okuda, S., Yoshizawa, A.C., and Kanehisa, M. (2007). KAAS: an automatic genome annotation and pathway reconstruction server. *Nucleic Acids Res.* *35*, W182–5.

59. Lowe, T.M., and Eddy, S.R. (1997). tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* *25*, 955–964.
60. Emanuelsson, O., Brunak, S., von Heijne, G., and Nielsen, H. (2007). Locating proteins in the cell using TargetP, SignalP and related tools. *Nat. Protoc.* *2*, 953–971.
61. Claros, M.G., and Vincens, P. (1996). Computational method to predict mitochondrially imported proteins and their targeting sequences. *Eur. J. Biochem.* *241*, 779–786.
62. Borgese, N., Colombo, S., and Pedrazzini, E. (2003). The tale of tail-anchored proteins: coming from the cytosol and looking for a membrane. *J. Cell Biol.* *161*, 1013–1019.
63. Imai, K., Fujita, N., Gromiha, M.M., and Horton, P. (2011). Eukaryote-wide sequence analysis of mitochondrial β -barrel outer membrane proteins. *BMC Genomics* *12*, 79.
64. Zubáčová, Z., Krylov, V., and Tachezy, J. (2011). Fluorescence in situ hybridization (FISH) mapping of single copy genes on *Trichomonas vaginalis* chromosomes. *Mol. Biochem. Parasitol.* *176*, 135–137.