


# Current Biology

## Phylogenomics Reveals Convergent Evolution of Lifestyles in Close Relatives of Animals and Fungi

### Graphical Abstract



### Authors

Guifré Torruella, Alex de Mendoza, Xavier Grau-Bové, ..., Ariadna Sitjà-Bobadilla, Stuart Donachie, Iñaki Ruiz-Trillo

### Correspondence

inaki.ruiz@ibe.upf-csic.es

### In Brief

Torruella et al. provide new molecular data from several protists and infer a novel phylogenomic framework for the opisthokonts that suggests rampant convergent evolution of several characters. Using comparative genomics, the authors show independent losses of the flagellum and delineate the evolutionary history of chitin synthases in this lineage.

### Highlights

- Taxon-rich phylogenomics provides an evolutionary framework for the opisthokonts
- Specialized osmotrophy evolved independently in fungi and animal relatives
- Opisthokonts underwent independent secondary losses of the flagellum
- The last opisthokont common ancestor had a complex repertoire of chitin synthases



# Phylogenomics Reveals Convergent Evolution of Lifestyles in Close Relatives of Animals and Fungi

Guifré Torruella,<sup>1,2,12</sup> Alex de Mendoza,<sup>1,2,12</sup> Xavier Grau-Bové,<sup>1,2</sup> Meritxell Antó,<sup>1</sup> Mark A. Chaplin,<sup>3</sup> Javier del Campo,<sup>1,4</sup> Laura Eme,<sup>5</sup> Gregorio Pérez-Cordón,<sup>6</sup> Christopher M. Whipps,<sup>7</sup> Krista M. Nichols,<sup>8,9</sup> Richard Paley,<sup>10</sup> Andrew J. Roger,<sup>5</sup> Ariadna Sitjà-Bobadilla,<sup>6</sup> Stuart Donachie,<sup>3</sup> and Iñaki Ruiz-Trillo<sup>1,2,11,\*</sup>

<sup>1</sup>Institut de Biologia Evolutiva (CSIC-Universitat Pompeu Fabra), Passeig Marítim de la Barceloneta 37-49, Barcelona 08003, Catalonia, Spain

<sup>2</sup>Departament de Genètica, Universitat de Barcelona, Avinguda Diagonal 645, Barcelona 08028, Catalonia, Spain

<sup>3</sup>Department of Microbiology, University of Hawaii at Manoa, Snyder Hall, 2538 McCarthy Mall, Honolulu, HI 96822, USA

<sup>4</sup>Department of Botany, University of British Columbia, Vancouver, BC V6T 1Z4, Canada

<sup>5</sup>Department of Biochemistry and Molecular Biology, Centre for Comparative Genomics and Evolutionary Bioinformatics, Dalhousie University, Halifax, NS B3H 4R2, Canada

<sup>6</sup>Institute of Aquaculture Torre de la Sal, IATS-CSIC, Ribera de Cabanes s/n, Castelló 12595, Spain

<sup>7</sup>Environmental and Forest Biology, State University of New York College of Environmental Science and Forestry (SUNY-ESF), Syracuse, NY 13210, USA

<sup>8</sup>Department of Biological Sciences, Purdue University, West Lafayette, IN 47907, USA

<sup>9</sup>Conservation Biology Division, Northwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, 2725 Montlake Boulevard East, Seattle, WA 98112, USA

<sup>10</sup>Centre for Environment Fisheries and Aquaculture Science, Weymouth Laboratory, Barrack Road, The Nothe, Weymouth, Dorset DT4 8UB, UK

<sup>11</sup>Institució Catalana de Recerca i Estudis Avançats (ICREA), Passeig Lluís Companys 23, Barcelona 08010, Catalonia, Spain

<sup>12</sup>Co-first author

\*Correspondence: [inaki.ruiz@ibe.upf-csic.es](mailto:inaki.ruiz@ibe.upf-csic.es)

<http://dx.doi.org/10.1016/j.cub.2015.07.053>

## SUMMARY

The Opisthokonta are a eukaryotic supergroup divided in two main lineages: animals and related protistan taxa, and fungi and their allies [1, 2]. There is a great diversity of lifestyles and morphologies among unicellular opisthokonts, from free-living phagotrophic flagellated bacterivores and filopodiated amoebas to cell-walled osmotrophic parasites and saprotrophs. However, these characteristics do not group into monophyletic assemblages, suggesting rampant convergent evolution within Opisthokonta. To test this hypothesis, we assembled a new phylogenomic dataset via sequencing 12 new strains of protists. Phylogenetic relationships among opisthokonts revealed independent origins of filopodiated amoebas in two lineages, one related to fungi and the other to animals. Moreover, we observed that specialized osmotrophic lifestyles evolved independently in fungi and protistan relatives of animals, indicating convergent evolution. We therefore analyzed the evolution of two key fungal characters in Opisthokonta, the flagellum and chitin synthases. Comparative analyses of the flagellar toolkit showed a previously unnoticed flagellar apparatus in two close relatives of animals, the filasterean *Ministeria vibrans* and *Corallochytrium limacisporum*. This implies that at least four different opisthokont lineages

secondarily underwent flagellar simplification. Analysis of the evolutionary history of chitin synthases revealed significant expansions in both animals and fungi, and also in the Ichthyosporea and *C. limacisporum*, a group of cell-walled animal relatives. This indicates that the last opisthokont common ancestor had a complex toolkit of chitin synthases that was differentially retained in extant lineages. Thus, our data provide evidence for convergent evolution of specialized lifestyles in close relatives of animals and fungi from a generalist ancestor.

## RESULTS AND DISCUSSION

### Broad Taxonomic Sampling Provides New Phylogenetic Insights into the Evolution of the Opisthokonta

Previous attempts to solve opisthokont phylogeny swayed between species-rich datasets with poor deep-node resolution based on small ribosomal subunit [1–3] and multigene supermatrices that included few taxa [4–6]. To improve upon our previously published phylogenomic dataset [6], we therefore sampled representative species in all described opisthokont lineages (see Table S1 and Supplemental Experimental Procedures). This included representatives of nucleariids, choanoflagellates, filastereans, and the two main lineages of Ichthyosporea (Dermocystidia and Ichthyophonida). In addition, we included two different strains of the enigmatic *Corallochytrium limacisporum*, a spherical free-living walled saprotroph found in coral reefs [7]. Originally classified as a thraustochytrid based on its morphology,

*C. limacisporum* has been unstably placed within the Opisthokonta in all molecular phylogenies to date because of the scarce molecular data available [8–11]. In order to improve the opisthokont outgroup, we also sampled the ancyromonad *Nutomonas longa* CCAP 1958/5 [12], which is putatively related to Apusomonadida [11]. Overall, we generated new transcriptomic data for 10 protistan taxa (11 strains in total, highlighted in bold in Figure 1), plus new genomic data from another strain (*Ichthyophonus hoferi*). This represents the broadest taxon sampling to date to infer the opisthokont phylogeny.

To investigate the phylogenetic relationships, we assembled two datasets comprising a total of 93 single-copy protein domains: one with 83 taxa and 18,218 aligned amino acid positions (S83), and the other with 70 taxa and 22,313 amino acid positions (S70). The latter dataset was constructed to maximize alignment length and to minimize topological artifacts by excluding putative problematic taxa with long branches (e.g., Microsporidia, Excavata) and high percentages of missing data (e.g., taxa with only expressed sequence tag data) (see Table S1). Both datasets were consistent in recovering the backbone of the eukaryotic phylogeny using both Bayesian inference (BI) (Figures 1 and S1C) and maximum likelihood (ML) (Figures S1A and S1B; see Supplemental Experimental Procedures for details).

As sister groups to Opisthokonta, we recovered Apusomonadida and Breviatea as recently reported [13], branching as independent lineages and not forming a monophyletic group or clustering with amoebozoans. Interestingly, the topology of the S83 dataset placed *Nutomonas longa* (Ancyromonadida) branching closer to the Excavata and not closely related to the Apusomonadida and Opisthokonta. This contrasts with previous analyses [11, 12] but is consistent with recent results based on multiple markers [14]. Within the Holomycota (which includes fungi and their protistan relatives), we recovered a clade formed by *Nuclearia* sp. and *Fonticula alba* (Discicristoidea) as the earliest-branching lineage [15]. This was followed by *Rozella alomycis* and Microsporidia [16] and the paraphyletic assemblage of Chytridiomycota (including Neocallimastigomycota) and Blastocladiomycota [17]. Finally, within the Holozoa we recovered Filasterea as the sister group to the clade formed by the Metazoa and Choanoflagellata, as previously reported [5, 6].

Interestingly, we recovered *C. limacisporum* as a sister group to Ichthyosporaea (including the two major groups Ichthyophonida and Dermocystida) [18] with both ML and BI methods. The S83 dataset recovered this position for *C. limacisporum* with weak support (56% ML bootstrap support [bs] and 0.8 BI posterior probability [pp]). However, support for this branch increased significantly (bs = 80%, pp = 0.84) when the long-branch taxa were excluded (see Figure 1 and Table S2). The position of the dermocystid *Sphaerothecum destruens* as sister group to the rest of ichthyosporaeans was only moderately supported (S83: bs = 60%, pp = 0.97; S70: bs = 61%, pp = 0.87) but was consistently recovered in all analyses. Thus, the monophyletic group comprising Ichthyosporaea and *C. limacisporum* appears to be the earliest-branching lineage in the Holozoa. We tentatively name this novel group “Teretosporea,” meaning “rounded spores,” through this study.

*C. limacisporum* is the only known free-living osmotroph in the Holozoa, whereas the ichthyosporaeans thus far described are known to be associated with animal hosts as parasites or com-

mensals [18], despite being frequently found in environmental surveys [3]. The life cycles of *C. limacisporum* and Ichthyosporaeans [7, 18] are strikingly similar: both start as a single cell that grows as a coenocyte until it reaches maturation, when it undergoes schizogony. The dispersive amoeboid or flagellated progeny (merozoites) settle and close the cycle [18]. Chytrid fungi show a similar developmental mode, with both coenocytic growth and amoeboid or flagellated stages [19]. Similarly, fungi also evolved from phagotrophic ancestors (Discicristoidea, *Rozella*, and Aphelida [20]) to become saprotrophs and parasites. Moreover, some Ichthyosporaea species (*A. parasiticum* and *I. hoferi*) present a mode of polar growth that clearly resembles fungal hyphae [21]. Thus, teretosporeans and fungi present tantalizing similarities regarding life style adaptations and morphologies.


The resulting opisthokont tree also confirms the convergent evolution of filose amoebas, Filasterea within the Holozoa and Discicristoidea within the Holomycota. Both lineages have evolved a similar cell morphology comprising long, actin-based filopodia [22], with some taxa going through an aggregative multicellular cell stage in their life cycles [23].

### Independent Loss of the Flagellum within the Opisthokonta

A single posterior motile flagellum is a defining character of opisthokonts [2]. Our observation that both filose amoebas and fungal-like lineages evolved in independent branches within opisthokonts therefore predicts independent loss of the flagellum. To address this hypothesis, we analyzed the evolution of the flagellar toolkit [24, 25]. The molecules that comprise the flagellum include specialized tubulins (*epsilon*, *delta*) [26], the intra-flagellar transport system (i.e., the IFT-A, IFT-B, and BBSome complexes [27]), and some motor molecules, mainly specialized subfamilies of dyneins and kinesins [24, 28] (Figure 2B). Large-scale genomic analyses have shown that the presence of these genes in a given genome correlates with the presence of a flagellum—revealing, in some cases, a previously unseen flagellar stage [28].


To clarify the evolution of the flagellum, we sought orthologs of a set of over 60 flagellum-specific proteins [24, 27, 28] in our taxon sampling (see Supplemental Experimental Procedures and Table S3). As expected, non-flagellated lineages such as Dikarya fungi, Discicristoidea, Ichthyophonida, and the filasterean *Capsaspora owczarzaki* yielded no significant hits (Figure 2A). This confirmed the recurrent secondary loss of the flagellum in at least four opisthokont lineages. In contrast, we found several proteins corresponding to key flagellar molecular components in the transcriptome of two taxa assumed not to be flagellated, the filasterean *M. vibrans* and the teretosporean *C. limacisporum*.

*M. vibrans* was originally described as a filose amoeba suspended in the water column by a stalk attached to the substrate. The stalk resembled a modified flagellum based on transmission electron microscopy (TEM) observations, which included structures resembling, according to the authors, doublet microtubules [2]. Interestingly, we observed the presence of axonemal dyneins, *epsilon* tubulin, and IFT-A/B complexes, clearly suggesting the presence of a flagellum in this species. Therefore, we tested whether the stalk is a modified flagellum by tubulin immunostaining on the original ATCC 50519 strain (see Supplemental Experimental Procedures). Confocal microscopy



**Figure 1. Phylogeny and Cell Biology of Opisthokonts**

(A) Phylogenetic tree based on the 83-taxa matrix (see Tables S1 and S2 and Supplemental Experimental Procedures) and inferred by PhyloBayes under the CAT-Poisson model. Tree topology is the consensus of two Markov chain Monte Carlo chains run for 1,500 generations, saving every ten trees and after a burn-in of 25%. Split supports are posterior probabilities (pp) and nonparametric maximum likelihood (ML) bootstrap (bs) values obtained from 200 ML replicates using the LG+I+G model implemented in RAXML. Support values > 0.95 pp and > 95% bs are indicated by a bullet (•). The taxa sampled in this study are indicated in bold. For raw trees, see Figure S1. (B–E) Light micrographs showing the coenocytic stage of representative species of the tentatively named "Teretosporea" (*Corallochytrium* + Ichthyosporea) sequenced in this study, including *Corallochytrium limacisporum* (B), *Sphaerothecum destruens* (C; arrowhead indicates flagellated zoospore), *Abeoforma whisleri* (D), and *Ichthyophonus hoferi* (E). Scale bar represents 10  $\mu$ m in (B)–(D) and 100  $\mu$ m in (E).



**Figure 2. Multiple Independent Losses of the Flagellar Toolkit and CHS Genes in Opisthokonta**

(A) Presence versus absence of key molecular components of the flagellar apparatus and chitin synthases (CHS) in distinct Opisthokonta lineages and taxa. <sup>a</sup>present in oomycetes, *Chlorella variabilis*, and *Paramecium tetraurelia*; <sup>b</sup>present in *Acanthamoeba castellanii*; <sup>c</sup>present in *Entamoeba histolytica* and *Thecamonas trahens*; <sup>d</sup>present in *Thalassiosira pseudonana*; <sup>e</sup>present in the chytrid *Batrachochytrium dendrobatidis*.

(B) Components of the flagellar apparatus and names of the molecular complexes. Adapted from [24]. See flagellar gene distribution in Table S3.

(C) Main chitin synthase classes and their canonical protein domain architectures (see CHS phylogeny in Figure S2).

revealed a tubulin protrusion branching from the cell body, which was specifically stained with  $\alpha$ -tubulin (Figure 3A) and acetylated tubulin antibodies (Figures 3B and S3). Moreover, our own TEM observations revealed a putative dense basal body and a flagellar section with nine outer ring structures and central microtubules (Figure 3C). Our transcriptomic data and experimental analysis thus revealed a flagellar structure in *M. vibrans*. Consequently, the ancestral filasterean must have had a flagellum, which was secondarily lost from *C. owczarzaki*.

The transcriptome of *C. limacisporum* was found to contain *delta/epsilon* tubulins, IFT-A and IFT-B components, and the retrograde motor kinesin-II (Figure 2A). Although this organism does possess an ortholog of HEATR2 recently linked to motile cilia [29], we did not find evidence of flagellar motility components, such as cytoplasmic dynein 2 or any of the axonemal dyneins (heavy, light, and intermediate chains; Table S3). Consistent with the original description of *C. limacisporum* [7], we did not observe a flagellum using light and TEM microscopy, at least under the culturing conditions employed. Therefore, our data suggest that *C. limacisporum* has a cryptic flagellated stage in its life cycle, as has been inferred for other eukaryotes (i.e., *Aureococcus* and *Ostreococcus*) based on their genome sequences [28]. Consequently, within the Teretosporea, a flagellated stage would be a feature shared by *C. limacisporum* and Dermocystida that was secondarily lost from the Ichthyophonida (Figure 4). This confirms the recurrent loss of the flagellum in both filose amoeboid lineages (Discicristoidea and Filasterea) and specialized osmotrophic lineages (Fungi and Teretosporea).


#### At Least Four Chitin Synthases in the Last Opisthokonta Common Ancestor

Given the apparent similarities in the evolution of the Fungi and Teretosporea, we investigated the evolutionary history of

another feature of fungal evolution, the cell wall. Chitin is a key biopolymer present in some fungal cell walls and animal cuticles [30], synthesized by chitin synthases (CHS), a large and complex multigene family. Several CHS classes have been described in fungi (classes I/II/III from division I and classes IV/V/VI/VII from division II) [31], with three ancestral classes known in animals [32]. Some fungal CHS classes are held as molecular synapomorphies of fungi (classes IV/V/VI/VII from division II), as they have been found exclusively in the genomes of fungi, including *R. allomycis* and microsporidian genomes [33]. Moreover, CHS homologs with uncertain classification have been found in other eukaryotes, including the oomycete *Saprolegnia monoica* [34], diatoms [35], and unicellular holozoans [18, 36].

To investigate which CHS classes are present in Teretosporea and to clarify their phylogenetic relationships with those in fungi and animals, we gathered CHS sequences from all eukaryotic supergroups and built a tree based on the chitin synthase domain (see Supplemental Experimental Procedures and Figure S2). This revealed three genes in *C. limacisporum* that belong to division II CHS and branch within the clade that comprises fungal classes IV/V/VII. These sequences consistently present the canonical functional motifs of fungal sequences (see Table S4). Interestingly, two of the genes encode an N-terminal myosin head domain, resembling genes from fungal classes V/VII [36] (Figure 2C). The myosin head of *C. limacisporum* CHS is sister group to fungal V/VII CHS, forming the myosin class XVII [37]. We thus propose that the CHS class IV/V/VII containing a myosin domain is an ancestral state in the Opisthokonta.


We also found that the Ichthyophonida contain CHS from both division I and division II clades. Ichthyophonida homologs from division I form a new clade with various eukaryotic sequences, including diatoms, choanoflagellates, and amoebozoans (Figures 2A and S2), revealing it also to be an ancestral class in



**Figure 3. Confocal and Electron Microscopy of *Ministeria vibrans* Flagellum**

(A and B) Confocal microscopy showing *Ministeria vibrans* ATCC 50519 stained with DAPI (blue) and anti- $\alpha$ -tubulin antibody 12G10 (Developmental Studies Hybridoma Bank) (green) (A) or with DAPI (blue), anti-acetylated-tubulin antibody T7451 (Sigma) (red), and phalloidin (green) (B). Arrowheads indicate the flagellar structure. Whereas the flagellar structure is specifically stained with cilia marker (acetylated tubulin) in (B), the cytoplasmic tubulin cytoskeleton is stained only with general anti-tubulin antibody in (A). *M. vibrans* feeds on bacteria, seen here as DAPI-stained bodies outside the cell. Scale bar represents 5  $\mu$ m. See also Figure S3.

(C) TEM micrograph showing a transverse section of the flagellar structure of *M. vibrans*. N, nucleus; F, flagellar structure; OM, outer microtubules; CM, central microtubules. Scale bar represents 200 nm.



**Figure 4. Evolution of Lifestyles and Some Cell Features of the Opisthokonta**

Opisthokonta cladogram displaying lifestyle characteristics such as feeding mode, flagellated stage, CHS repertoire, and developmental mode (see Figure S4 for wheat germ agglutinin [WGA] staining) and ancestral state reconstruction of the last opisthokont common ancestor (LOCA). Choanoflagellate image is adapted from <http://www.dayel.com/> (CC BY-SA 3.0).

the eukaryotes. Ichthyosporean division II CHS homologs belong to the Metazoan class, which is also present in other unicellular holozoans, apusomonads, and amoebozoans but is secondarily lost in fungi. Finally, fungal class I/II/III is found in several bikonta, including oomycetes and chlorophytes, suggesting an ancestral origin and secondary loss from the Holozoa. In summary, at least four ancestral paralogs of structurally different CHS (Figure 2C) were found in the last opisthokont common ancestor (LOCA), and secondary loss appears to have been common in descendant lineages (Figure 4). The presence of a complex CHS repertoire in the ancestor of all Opisthokonta, and the retention of rich CHS repertoires in the cell-walled lineages, suggests that the presence of chitin in the cell wall was an ancestral feature and not a fungal synapomorphy [33]. Consistent with this suggestion, Ichthyosporeans encoding a complex CHS repertoire showed chitin staining in the cell wall (Figure S4), and therefore only CHS VI class and the diversification of CHS IV/V/VII class into paralogous groups could be still considered fungal molecular synapomorphies.

#### A New Phylogenetic Framework for the Opisthokonta

By obtaining the transcriptomes of 10 new protist taxa (11 strains), plus the genome of an additional strain (12 strains in total), we have improved the previously biased representation

of genomic information for unicellular Opisthokonta. This allowed us to reassess the phylogenetic relationships among the opisthokonts through an unprecedented gene- and taxon-rich approach. Our dataset, with few missing data (Table S1), includes representatives from all opisthokont lineages, providing a stronger phylogenetic framework for internal relationships. Our phylogenetic analyses reveal a new clade: [Ichthyosporia + *C. limacisporum*], which we tentatively call Teretosporea, and which represents the earliest holozoan divergence (Figure 1).

Our data reveal that convergent evolution explains similarities in the lifestyles of the Fungi and Teretosporea as well as in Filasterea and Discicristoidea (Figure 4). The ancestral LOCA was most likely a filopodiated and flagellated generalist bacterivore [38]. Consequently, the specialized osmotrophic feeding mode, cell wall, and transition from saprotrophic to parasitic lifestyles in Fungi and Teretosporea occurred independently. This is not rare in eukaryotes, since similar adaptations are also found in stramenopiles such as the oomycetes and the thraustochytrids [39, 40]. However, our data provide the first example of such a process occurring in a close relative of animals. Through analysis of secondary loss of the flagellum and differential retention of ancestral CHS paralogs in opisthokonts, we have also provided molecular evidence to explain these lifestyle adaptations. Therefore, this study provides a striking example of convergent evolution through differential retention of ancestral genomic characters in the unicellular relatives of animals and fungi.

#### ACCESSION NUMBERS

The accession numbers for new data reported in this study are NCBI Sequence Read Archive: SRS502375, SRS502376, SRS721318, SRS725979, SRS725801, SRS726091, SRS724896, SRS725006, SRX179384; and NCBI BioProject: PRJNA290639.

#### SUPPLEMENTAL INFORMATION

Supplemental Information includes four figures, four tables, and Supplemental Experimental Procedures and can be found with this article online at <http://dx.doi.org/10.1016/j.cub.2015.07.053>.

#### AUTHOR CONTRIBUTIONS

G.T., A.d.M., and I.R.-T. designed and coordinated the study. *C. limacisporum* isolation was performed by M.A.C. and S.D. *I. hoferi* genomic data was performed by K.M.N. and C.M.W. *S. destruens* RNA data was performed by R.P. Other strain cultivation and RNA extractions were performed by G.T., J.d.C., M.A., G.P.-C., and X.G.-B. Phylogenomics was performed by G.T., L.E., and A.J.R. Flagellum and CHS comparative genomics were performed by A.d.M. WGA staining was performed by A.d.M. and M.A. Immunostaining and TEM were performed by X.G.-B., M.A., A.S.-B., and G.T. Figures were assembled by X.G.-B., G.T., and A.d.M. G.T., A.d.M., and I.R.-T. wrote the manuscript. All authors commented on the manuscript.

#### ACKNOWLEDGMENTS

This work was supported by two grants (BFU2011-23434 and BFU2014-57779-P) from Ministerio de Economía y Competitividad (MINECO) and an ERC Starting Grant (ERC-2007-StG-206883) to I.R.-T. We also acknowledge support from Secretaria d'Universitats i Recerca del Departament d'Economia i Coneixement de la Generalitat de Catalunya (project 2014 SGR 619). G.T. was supported by a pre-graduate FI grant from the Catalan Government. A.d.M. and X.G.-B. were supported by a pre-graduate FPI grant from MINECO. L.E. was supported by a CGEB postdoctoral fellowship from

the Tula Foundation, and A.J.R. was supported by a CIHR-NSHRF RPP grant (FRN# 62809). Assembly of the *I. hoferi* genome (USA) was supported by NSF grant number ACI-1053575. G.P.-C. was supported by a "Juan de la Cierva" grant, and A.S.-B. was supported by Generalitat Valenciana (PROMETEO FASE II-2014/085). We thank Dan Richter, Nicole King, Franz Lang, Matt Brown, Joseph Ryan, Andy Baxeavanis, Ana Riesgo, Scott Nichols, Romain Derelle, Jordi Paps, Diego Mallo, Martin Kolisko, Txema Heredia, Philippe Lopez, Eric Baptiste, Arnau Sebé-Pedrós, Ana Franco-Sierra, and Jake L. Gregg for providing samples, data, technical assistance, and/or insightful comments.

Received: November 10, 2014

Revised: June 30, 2015

Accepted: July 22, 2015

Published: September 10, 2015

#### REFERENCES

- Medina, M., Collins, A.G., Taylor, J.W., Valentine, J.W., Lipps, J.H., Amaral-Zettler, L., and Sogin, M.L. (2003). Phylogeny of Opisthokonta and the evolution of multicellularity and complexity in Fungi and Metazoa. *Int. J. Astrobiol.* 2, 203–211.
- Cavalier-Smith, T., and Chao, E.E.-Y. (2003). Phylogeny of choanozoa, apusozoa, and other protozoa and early eukaryote megaevolution. *J. Mol. Evol.* 56, 540–563.
- del Campo, J., and Ruiz-Trillo, I. (2013). Environmental survey meta-analysis reveals hidden diversity among unicellular opisthokonts. *Mol. Biol. Evol.* 30, 802–805.
- Ruiz-Trillo, I., Roger, A.J., Burger, G., Gray, M.W., and Lang, B.F. (2008). A phylogenomic investigation into the origin of metazoa. *Mol. Biol. Evol.* 25, 664–672.
- Shalchian-Tabrizi, K., Minge, M.A., Espelund, M., Orr, R., Ruden, T., Jakobsen, K.S., and Cavalier-Smith, T. (2008). Multigene phylogeny of choanozoa and the origin of animals. *PLoS ONE* 3, e2098.
- Toruella, G., Derelle, R., Paps, J., Lang, B.F., Roger, A.J., Shalchian-Tabrizi, K., and Ruiz-Trillo, I. (2012). Phylogenetic relationships within the Opisthokonta based on phylogenomic analyses of conserved single-copy protein domains. *Mol. Biol. Evol.* 29, 531–544.
- Raghukumar, S. (1987). Occurrence of the thraustochytrid, *Corallochytrium limacisporum* gen. et sp. nov. in the coral reef lagoons of the Lakshadweep islands in the Arabian Sea. *Bot. Mar.* 30, 83–89.
- Ruiz-Trillo, I., Lane, C.E., Archibald, J.M., and Roger, A.J. (2006). Insights into the evolutionary origin and genome architecture of the unicellular opisthokonts *Capsaspora owczarzaki* and *Sphaeroforma arctica*. *J. Eukaryot. Microbiol.* 53, 379–384.
- Steenkamp, E.T., Wright, J., and Baldauf, S.L. (2006). The protistan origins of animals and fungi. *Mol. Biol. Evol.* 23, 93–106.
- Sumathi, J.C., Raghukumar, S., Kasbekar, D.P., and Raghukumar, C. (2006). Molecular evidence of fungal signatures in the marine protist *Corallochytrium limacisporum* and its implications in the evolution of animals and fungi. *Protist* 157, 363–376.
- Paps, J., Medina-Chacón, L.A., Marshall, W., Suga, H., and Ruiz-Trillo, I. (2013). Molecular phylogeny of unikonts: new insights into the position of apusomonads and ancyromonads and the internal relationships of opisthokonts. *Protist* 164, 2–12.
- Glücksman, E., Snell, E.A., and Cavalier-Smith, T. (2013). Phylogeny and evolution of Planomonadida (Sulcozoa): eight new species and new genera *Fabomonas* and *Nutomonas*. *Eur. J. Protistol.* 49, 179–200.
- Brown, M.W., Sharpe, S.C., Silberman, J.D., Heiss, A.A., Lang, B.F., Simpson, A.G., and Roger, A.J. (2013). Phylogenomics demonstrates that breviate flagellates are related to opisthokonts and apusomonads. *Proc. Biol. Sci.* 280, 20131755.
- Cavalier-Smith, T., Chao, E.E., Snell, E.A., Berney, C., Fiore-Donno, A.M., and Lewis, R. (2014). Multigene eukaryote phylogeny reveals the likely protozoan ancestors of opisthokonts (animals, fungi, choanozoans) and Amoebozoa. *Mol. Phylogenet. Evol.* 81, 71–85.

15. Liu, Y., Steenkamp, E.T., Brinkmann, H., Forget, L., Philippe, H., and Lang, B.F. (2009). Phylogenomic analyses predict sistergroup relationship of nucleariids and fungi and paraphyly of zygomycetes with significant support. *BMC Evol. Biol.* **9**, 272.
16. James, T.Y., Letcher, P.M., Longcore, J.E., Mozley-Standridge, S.E., Porter, D., Powell, M.J., Griffith, G.W., and Vilgalys, R. (2006). A molecular phylogeny of the flagellated fungi (Chytridiomycota) and description of a new phylum (Blastocladiomycota). *Mycologia* **98**, 860–871.
17. James, T.Y., Kauff, F., Schoch, C.L., Matheny, P.B., Hofstetter, V., Cox, C.J., Celio, G., Gueidan, C., Fraker, E., Miądlikowska, J., et al. (2006). Reconstructing the early evolution of Fungi using a six-gene phylogeny. *Nature* **443**, 818–822.
18. Gockling, S.L., Marshall, W.L., and Gleason, F.H. (2013). Phylogenetic interpretations and ecological potentials of the Mesomycetozoa (Ichthyosporia). *Fungal Ecol.* **6**, 237–247.
19. Shelest, K., and Voigt, K. (2014). Genomics to study basal lineage fungal biology: Phylogenomics suggests a common origin. In *Fungal Genomics*, M. Nowrousian, ed. (Springer-Verlag), pp. 31–60.
20. Karpov, S.A., Mamkaeva, M.A., Aleoshin, V.V., Nassonova, E., Lilje, O., and Gleason, F.H. (2014). Morphology, phylogeny, and ecology of the apheleids (Aphelidea, Opisthokonta) and proposal for the new superphylum Opisthosporidia. *Front. Microbiol.* **5**, 112.
21. Gozlan, R.E., Marshall, W.L., Lilje, O., Jessop, C.N., Gleason, F.H., and Andreou, D. (2014). Current ecological understanding of fungal-like pathogens of fish: what lies beneath? *Front. Microbiol.* **5**, 62.
22. Sebé-Pedrós, A., Burkhardt, P., Sánchez-Pons, N., Fairclough, S.R., Lang, B.F., King, N., and Ruiz-Trillo, I. (2013). Insights into the origin of metazoan filopodia and microvilli. *Mol. Biol. Evol.* **30**, 2013–2023.
23. Sebé-Pedrós, A., Irimia, M., Del Campo, J., Parra-Acero, H., Russ, C., Nusbaum, C., Blencowe, B.J., and Ruiz-Trillo, I. (2013). Regulated aggregative multicellularity in a close unicellular relative of metazoa. *eLife* **2**, e01287.
24. Carvalho-Santos, Z., Azimzadeh, J., Pereira-Leal, J.B., and Bettencourt-Dias, M. (2011). Evolution: Tracing the origins of centrioles, cilia, and flagella. *J. Cell Biol.* **194**, 165–175.
25. Hodges, M.E., Scheumann, N., Wickstead, B., Langdale, J.A., and Gull, K. (2010). Reconstructing the evolutionary history of the centriole from protein components. *J. Cell Sci.* **123**, 1407–1413.
26. Findeisen, P., Mühlhausen, S., Dempewolf, S., Hertzog, J., Zietlow, A., Carlomagno, T., and Kollmar, M. (2014). Six subgroups and extensive recent duplications characterize the evolution of the eukaryotic tubulin protein family. *Genome Biol. Evol.* **6**, 2274–2288.
27. van Dam, T.J.P., Townsend, M.J., Turk, M., Schlessinger, A., Sali, A., Field, M.C., and Huynen, M.A. (2013). Evolution of modular intraflagellar transport from a coatomer-like progenitor. *Proc. Natl. Acad. Sci. USA* **110**, 6943–6948.
28. Wickstead, B., and Gull, K. (2012). Evolutionary biology of dyneins. In *Dyneins*, S. King, ed. (Elsevier), pp. 89–121.
29. Diggle, C.P., Moore, D.J., Mali, G., zur Lage, P., Ait-Lounis, A., Schmidts, M., Shoemark, A., Garcia Munoz, A., Halachev, M.R., Gautier, P., et al. (2014). *HEATR2* plays a conserved role in assembly of the ciliary motile apparatus. *PLoS Genet.* **10**, e1004577.
30. Merzendorfer, H. (2011). The cellular basis of chitin synthesis in fungi and insects: common principles and differences. *Eur. J. Cell Biol.* **90**, 759–769.
31. Ruiz-Herrera, J., and Ortiz-Castellanos, L. (2010). Analysis of the phylogenetic relationships and evolution of the cell walls from yeasts and fungi. *FEMS Yeast Res.* **10**, 225–243.
32. Zakrzewski, A.-C., Weigert, A., Helm, C., Adamski, M., Adamska, M., Bleidorn, C., Raible, F., and Hausen, H. (2014). Early divergence, broad distribution, and high diversity of animal chitin synthases. *Genome Biol. Evol.* **6**, 316–325.
33. James, T.Y., Pelin, A., Bonen, L., Ahrendt, S., Sain, D., Corradi, N., and Stajich, J.E. (2013). Shared signatures of parasitism and phylogenomics unite Cryptomycota and microsporidia. *Curr. Biol.* **23**, 1548–1553.
34. Leal-Morales, C.A., Gay, L., Fèvre, M., and Bartnicki-García, S. (1997). The properties and localization of *Saprolegnia monoica* chitin synthase differ from those of other fungi. *Microbiology* **143**, 2473–2483.
35. Durkin, C.A., Mock, T., and Armbrust, E.V. (2009). Chitin in diatoms and its association with the cell wall. *Eukaryot. Cell* **8**, 1038–1050.
36. James, T.Y., and Berbee, M.L. (2012). No jacket required—new fungal lineage defies dress code: recently described zoosporic fungi lack a cell wall during trophic phase. *BioEssays* **34**, 94–102.
37. Sebé-Pedrós, A., Grau-Bové, X., Richards, T.A., and Ruiz-Trillo, I. (2014). Evolution and classification of myosins, a paneukaryotic whole-genome approach. *Genome Biol. Evol.* **6**, 290–305.
38. Cavalier-Smith, T. (2013). Early evolution of eukaryote feeding modes, cell structural diversity, and classification of the protozoan phyla Loukozoa, Sulcozoa, and Choanozoa. *Eur. J. Protistol.* **49**, 115–178.
39. Spanu, P., and Kämper, J. (2010). Genomics of biotrophy in fungi and oomycetes—emerging patterns. *Curr. Opin. Plant Biol.* **13**, 409–414.
40. Richards, T.A., and Talbot, N.J. (2013). Horizontal gene transfer in osmotrophs: playing with public goods. *Nat. Rev. Microbiol.* **11**, 720–727.